

Susac syndrome

Version: 3.0

Published: 20 October 2018

Condition for which IVIg use is in exceptional circumstances only

Specific Conditions	<ul style="list-style-type: none">• Susac syndrome
Indication for IVIg Use	<ul style="list-style-type: none">• Probable or definite Susac syndrome in concurrence with high dose corticosteroids
Level of Evidence	Insufficient data (Category 4a)
Description and Diagnostic Criteria	Susac syndrome is a rare, microangiopathic disorder characterised by encephalopathy, hearing loss and retinal artery branch occlusions. Case reports show benefit of intravenous immunoglobulin (IVIg) or plasma exchange in combination with corticosteroids, generally with, or in mild cases without, other immunosuppressive agents.
Justification for Evidence Category	There are no randomised controlled trials (RCTs) nor prospective series in Susac syndrome. However, there is a very poor natural history and a clear response to multi-agent immunosuppressive therapy including intravenous immunoglobulin (IVIg) in case series (Mateen, 2012). A recent series has suggested equal or possibly greater efficacy from plasma exchange over IVIg, and plasma exchange should be considered where available (Vodopivec, 2016).
Diagnosis Requirements	A diagnosis must be made by an Immunologist, Neurologist, Rheumatologist or an Ophthalmologist.

Qualifying Criteria for IVIg Therapy

- Probable or definite diagnosis of Susac syndrome has been made by the presence of at least two of the following:
 - Encephalopathy with typical MRI brain changes including corpus callosum lesions or characteristic diffusion weighted imaging (DWI) hyperintense lesions
 - New sensorineural hearing loss or tinnitus
 - Branch retinal artery occlusions (BRAOs) or ischaemia or arterial wall hyperfluorescence (AWH) on fluorescein angiography

AND

- A baseline assessment of disability is conducted as measured by the [Modified Rankin Scale \(MRS\)](#)

AND

- Corticosteroid therapy is being given concurrently

OR

- Steroid therapy is absolutely contraindicated

IVIg should be used for four months (induction plus three maintenance cycles) before determining whether the patient has responded. If there is no benefit after this treatment, IVIg therapy should be abandoned.

Review by a neurologist, immunologist rheumatologist or ophthalmologist is required within four months of treatment to determine whether the patient has responded, and annually thereafter.

For stable patients on maintenance treatment, review by a neurologist, immunologist, ophthalmologist or rheumatologist is required at least annually. A trial off IVIg should be attempted after a year of therapy, unless there is a contraindication to doing so, or the patient has previously relapsed after an earlier trial of withdrawal of IVIg.

Documentation of clinical effectiveness is necessary for continuation of IVIg therapy.

Review Criteria for Assessing the Effectiveness of IVIg Use

IVIg should be used for four months (induction plus three maintenance cycles) before determining whether the patient has responded. If there is no benefit after this treatment, IVIg therapy should be abandoned.

Review by an neurologist, immunologist rheumatologist or ophthalmologist is required within four months of treatment to determine whether the patient has responded, and annually thereafter.

For stable patients on maintenance treatment, review by a neurologist, immunologist, ophthalmologist or rheumatologist is required at least annually. A trial off IVIg should be attempted after a year of therapy, unless there is a contraindication to doing so, or the patient has previously relapsed after an earlier trial of withdrawal of IVIg.

Documentation of clinical effectiveness is necessary for continuation of IVIg therapy.

Clinical effectiveness of Ig therapy can be demonstrated by:

On review of the initial authorisation period

- Stabilisation of, or improvement in the severity of symptoms
- AND
- Stabilisation of, or improvement in disability as measured by the [Modified Rankin Scale \(MRS\)](#) score compared to the qualifying assessment

On review of a continuing authorisation period

- Improvement in or stabilisation of symptoms compared to the previous review assessment
- AND
- Stabilisation or improvement in disability as measured by the [Modified Rankin Scale \(MRS\)](#) compared to the previous review assessment
- AND
- A trial of weaning towards cessation of Ig therapy is planned once the patient is stable or a reason is provided as to why a trial is not planned

A trial off IVIg should be attempted after a year of therapy, unless there is a contraindication to doing so, or the patient has previously relapsed after an earlier trial of withdrawal of IVIg.

Dose

- **Induction Dose** - Up to 2 g/kg over 2 to 5 days.
- **Maintenance Dose** - 0.5 - 1 g/kg every 2 to 6 weeks.

The aim should be to use the lowest dose possible that achieves the appropriate clinical outcome for each patient.

Refer to the current product information sheet for further information on dose, administration and contraindications.

Bibliography

Aubart-Cohen, F, Klein, I, Alexandra, JF, et al 2007, 'Long-term outcome in Susac syndrome', *Medicine (Baltimore)*, vol. 86, no. 2, pp. 93–102. <https://www.ncbi.nlm.nih.gov/pubmed/17435589>

Bonita, R & Beaglehole, R 1988, 'Recovery of motor function after stroke', *Stroke*, vol. 19, no. 12, pp. 1497-1500.

Bibliography

<https://www.ahajournals.org/doi/abs/10.1161/str.19.12.3201508>

Fox, RJ, Costello, F, Judkin, A, et al 2006, 'Treatment of Susac syndrome with gamma globulin and corticosteroids', *Journal of the Neurological Sciences*, vol. 251, no. 1–2, pp. 17–22. <https://www.ncbi.nlm.nih.gov/pubmed/17052732>

Mateen, FJ, Zubkov, AY, Muralidharan, R, et al 2012, 'Susac syndrome: clinical characteristics and treatment in 29 new cases', *European Journal of Neurology*, vol. 19, no. 6, pp. 800-11. <https://www.ncbi.nlm.nih.gov/pubmed/22221557>

Rankin, J, 1957, 'Cerebral vascular accidents in patients over the age of 60', *Scottish Medical Journal*, no. 2, pp. 200-15.

Stroke Society of Australia, *The Modified Rankin Scale*, Available from: http://www.strokesociety.com.au/index.php?option=com_content&view=article&id=292:modified-rankin-scale-astn&catid=40:astn

Van Swieten, JC, Koudstaal, PJ, Visser, MC, Schouten, HJ, van Gijn, J, 1987, 'Interobserver agreement for the assessment of handicap in stroke patients', *Stroke*, vol. 19, no. 5, pp. 604-607.
<https://pdfs.semanticscholar.org/fba2/da6e4888b08ee79441937169c53f16aec218.pdf>

Vodopivec, I & Prasad, S 2016, 'Treatment of Susac Syndrome', *Current Treatment Options in Neurology*, vol. 18, no. 1, pp. 3. <https://www.ncbi.nlm.nih.gov/pubmed/26715396>

Generated on: 7 April 2019