

Epidermolysis bullosa acquisita

Version: 3.0

Published: 20 October 2018

Condition for which IVIg use is in exceptional circumstances only

Specific Conditions	<ul style="list-style-type: none">• Epidermolysis bullosa acquisita
Indication for IVIg Use	<ul style="list-style-type: none">• Persistent severe EBA refractory to conventional immunosuppressive therapy• Treatment of an ongoing flare of EBA disease in responding patients who have ceased Ig therapy
Level of Evidence	Insufficient data (Category 4a)
Description and Diagnostic Criteria	<p>Epidermolysis bullosa acquisita (EBA) is a rare, potentially severe life-threatening disease which has no cure. This subepidermal blistering disorder of the skin and mucous membranes primarily affects adults but has been described in a few children. Although the majority of patients may have mild disease with blistering over trauma prone areas, there have been cases of severe widespread disease that remains refractory to conventional therapies.</p> <p>The end stage results of severe disease include ocular disease with conjunctival scarring and blindness, mucosal disease with oesophageal strictures and scarring fibrosis of the skin with alopecia, nail loss and mitten like deformities of the hands.</p> <p>EBA is an autoimmune disease characterised by the production of antibodies against type VII collagen resulting in immune-mediated disruption of the anchoring fibrils that connect the basement membrane to dermal structures and the clinical development of blistering.</p> <p>Diagnosis of EBA is based on history, full skin examination, and skin biopsies.</p> <p>There is limited data on treatment options for EBA and optimal approach to treatment has not been established. Suggested initial treatment is with colchicine or dapsone (Grade 2C). If treatment is not effective, these agents may be used simultaneously.</p> <p>EBA that is refractory to the above requires more aggressive therapy. Agents that may have efficacy for refractory EBA include immunosuppressants, intravenous immunoglobulin, and rituximab.</p>
Justification for Evidence Category	<p>Intravenous immunoglobulin (IVIg) may be an option in severe disease. Multiple case reports suggest that the periodic administration of IVIg alone or in combination with other agents is also effective for improving the clinical manifestations of epidermolysis bullosa acquisita (EBA). In a 2011 review of published reports, 14 of 15 patients, mostly with severe widespread refractory disease, given IVIg as monotherapy or in conjunction with other therapies achieved clinical improvement (Gurcan, 2011). Multiple cycles of IVIg were typically given; each cycle usually consisted of a total of 1.5 to 2g/kg of IVIg given over the course of three to five days. Thirteen of 15 patients remained on IVIg for maintenance treatment.</p> <p>A more recent retrospective case series demonstrated similar clinical response rates, but suggested IVIg may induce a more sustained remission (Ahmed, 2012). Ten patients, all nonresponsive to conventional treatments, were started on 2g/kg/cycle of IVIg for a mean of 23 cycles over 39 months. All 10 demonstrated clinical response and were able to completely withdraw their previous therapy; no recurrence was observed during a mean follow-up period of 54 months after cessation of treatment.</p>

Diagnosis Requirements	A diagnosis must be made by an Immunologist or a Dermatologist.
Qualifying Criteria for IVIg Therapy	<div>Persistent severe EBA refractory to conventional immunosuppressive therapy</div> <div> <ul style="list-style-type: none"> Persistent severe EBA disease confirmed by biopsy and/or immunofluorescence including ophthalmological and/or mucosal sites <p>AND</p> <ul style="list-style-type: none"> Persistent disease despite standard treatment with colchicine and dapsone and at least two other immunosuppressant agents <p>OR</p> <ul style="list-style-type: none"> Corticosteroid and/or immunosuppressant therapy is contraindicated or has resulted in unacceptable side effects or significant toxicity </div> <div>Treatment of an ongoing flare of EBA disease in responding patients who have ceased Ig therapy</div> <div> <ul style="list-style-type: none"> Ongoing flare of mucosal or ophthalmic EBA disease in patients with EBA disease confirmed by biopsy and/or immunofluorescence including ophthalmological and/or mucosal site <p>AND</p> <ul style="list-style-type: none"> Reduction in severity and/or the number of lesions was demonstrated in response to initial Ig therapy <p>AND</p> <ul style="list-style-type: none"> At least one immunosuppressant medication is to be given concurrently <p>IVIg should be used for four months (induction and three maintenance cycles) before determining whether the patient has responded. If the patient has not responded after this time, Ig therapy should be abandoned. Review is required by a dermatologist or immunologist after the first four months treatment to confirm response, and six monthly thereafter.</p> <p>Documentation of clinical effectiveness is necessary for continuation of IVIg therapy.</p> </div>

Review Criteria for Assessing the Effectiveness of IVIg Use

Persistent severe EBA refractory to conventional immunosuppressive therapy

Review is not mandated for this indication however the following criteria may be useful in assessing the effectiveness of Ig therapy.

- Reduction in number of blisters/erosions and improved healing compared to the level at the qualifying assessment

Treatment of an ongoing flare of EBA disease in responding patients who have ceased Ig therapy

IVIg should be used for four months (induction and three maintenance cycles) before determining whether the patient has responded. If the patient has not responded after this time, Ig therapy should be abandoned.

Review is required by a dermatologist or immunologist after the first four months treatment to confirm response, and six monthly thereafter.

Documentation of clinical effectiveness is necessary for continuation of IVIg therapy.

Clinical effectiveness of Ig therapy can be assessed by:

On review of the initial authorisation period

- Reduction in severity and number of erosions or blisters and improved healing compared to the level at the qualifying assessment

AND

- At least one immunosuppressant medication is given concurrently

On review of a continuing authorisation period

For stable patients on maintenance treatment, review by dermatologist or an immunologist is required six monthly.

Consideration should be given to a trial-off immunoglobulin (Ig) therapy once the patient has achieved stabilised disease or clinical remission.

Clinical effectiveness of Ig therapy may be assessed by:

- Reduction in the number of erosions or blisters and improved healing compared to the previous review

AND

- There is remaining activity or stable disease requiring further treatment

AND

- Immunosuppressant medication is given concurrently

AND

- A trial-off Ig therapy is planned or, if not planned, a reason is provided

Dose

Persistent severe EBA refractory to conventional immunosuppressive therapy

- **Initial therapy** - 1.5 – 2 g/kg over 3 to 5 days, monthly for three months.

Treatment should be for no longer than three months initially after which time a clinical response should be demonstrated, and the patient trialled off therapy. If the patient has not responded within this time, Ig therapy should be abandoned. If disease flares following cessation in responding patients, a request for low dose maintenance therapy can be made.

The aim should be to use the lowest dose possible that achieves the appropriate clinical outcome for each patient

Refer to the current product information sheet for further information on dose, administration and contraindications.

Treatment of an ongoing flare of EBA disease in responding patients who have ceased Ig therapy

- **Induction Dose** - 1.5 to 2 g/kg over 3 to 5 days.
- **Maintenance Dose** - 0.4 g/kg four to six weekly.

The aim should be to use the lowest dose possible that achieves the appropriate clinical outcome for each patient.

Refer to the current product information sheet for further information on dose, administration and contraindications.

Bibliography

Ahmed, AR & Gürcan, HM, 2012, 'Treatment of epidermolysis bullosa acquisita with intravenous immunoglobulin in patients non-responsive to conventional therapy: clinical outcome and post-treatment long-term follow-up', *Journal of the European Academy of Dermatology and Venerology*, vol. 26, no. 9, pp.1074-83.

<https://www.ncbi.nlm.nih.gov/pubmed/21819451>

Caldwell, JB, Yancey, KB, Engler, RJ, et al 1994, 'Epidermolysis bullosa acquisita: efficacy of high-dose intravenous immunoglobulins', *Journal of the American Academy of Dermatology*, vol. 31, pp. 827-8.

<https://www.ncbi.nlm.nih.gov/pubmed/7929940>

Gürcan, HM & Ahmed, AR, 2011, 'Current concepts in the treatment of epidermolysis bullosa acquisita', *Expert Opinion on Pharmacotherapy*, vol. 12, no. 8, pp. 1259-68. <https://www.ncbi.nlm.nih.gov/pubmed/21254861>

Iranzo, P, Herrero-González, JE & Mascaró-Galy, JM, et al 2014, 'Epidermolysis bullosa acquisita: a retrospective analysis of 12 patients evaluated in four tertiary hospitals in Spain', *British Journal of Dermatology*, vol. 171, no. 5, pp. 1022-30.

<https://www.ncbi.nlm.nih.gov/pubmed/24890437>

Kirtschig, G, Murrell, D, Wojnarowska, F, et al 2003, 'Interventions for mucous membrane pemphigoid and epidermolysis bullosa acquisita', *Cochrane Database of Systematic Reviews*, vol. 2003, no.1. pp. CD004056.

<https://www.ncbi.nlm.nih.gov/pubmed/12535507>

Kofler, H, Wambacher-Gasser, B, Topar, G, et al 1997, 'Intravenous immunoglobulin treatment in therapy-resistant epidermolysis bullosa acquisita', *Journal of the American Academy of Dermatology*, vol. 36, pp. 331-5.

<https://www.ncbi.nlm.nih.gov/pubmed/9039213>

Gourgiotou, K, Exadaktylou, D, Aroni, K, et al 2002, 'Epidermolysis bullosa acquisita: treatment with intravenous immunoglobulins', *Journal of the European Academy of Dermatology and Venerology*, vol. 16, no. 1, pp.77-80.

<https://www.ncbi.nlm.nih.gov/pubmed/11952298>

Bibliography

Meier, F, Sönnichsen, K, Schaumburg-Lever, G, et al 1993, 'Epidermolysis bullosa acquisita: efficacy of high-dose intravenous immunoglobulins', *Journal of the American Academy of Dermatology*, vol. 29, pp. 334-7.
<https://www.ncbi.nlm.nih.gov/pubmed/8340508>

Mohr, C, Sunderkötter, C, Hildebrand, A, et al 1995, 'Successful treatment of epidermolysis bullosa acquisita using intravenous immunoglobulins', *British Journal of Dermatology*, vol. 132, no. 5, pp. 824-6.
<https://www.ncbi.nlm.nih.gov/pubmed/7772494>

Oktem, A, Akay, BN, Boyvat, A, et al 2017, 'Long-term results of rituximab-intravenous immunoglobulin combination therapy in patients with epidermolysis bullosa acquisita resistant to conventional therapy', *Journal of Dermatological Treatment*, vol. 28, no. 1, pp.50-54. <https://www.ncbi.nlm.nih.gov/pubmed/27161164>

Ontario Regional Blood Coordinating Network 2016, 'Ontario Intravenous Immune Globulin (IVIG) Utilization Management Guidelines, Version 3.0. [online]'

UK Department of Health 2011, 'Clinical Guidelines for Immunoglobulin Use: Second Edition Update', Available from: <https://www.gov.uk/government/publications/clinical-guidelines-for-immunoglobulin-use-second-edition-update>

UK Department of Health 2011, 'Clinical Guidelines for Immunoglobulin Use: Second Edition Update: Summary Poster', Available from: <http://igd.mdsas.com/clinical-info/>

Generated on: 4 April 2019