

Sjögren’s syndrome associated neuropathy

Condition for which Ig use is in exceptional circumstances only

Specific Conditions	<ul style="list-style-type: none">Painful small fibre neuropathyAtaxic sensory neuronopathySensorimotor axonal neuropathyAutonomic neuropathy
Indication for Ig Use	<ul style="list-style-type: none">Severe, primary Sjögren’s syndrome associated neuropathy that is unresponsive to corticosteroid and immunosuppressant therapyRelapse of Sjögren’s syndrome associated neuropathy within six months of trial off Ig therapy
Level of Evidence	Insufficient data (Category 4a)
Description and Diagnostic Criteria	The diagnosis of primary Sjögren’s syndrome should be consistent with the criteria of the American-European Study group (Tzioufas et al, 2007). Intravenous immunoglobulin (IVIg) may be indicated in patients with some neuropathy subtypes associated with Sjögren’s syndrome (excluding those caused by necrotising vasculitis) where other treatments have been ineffective.
Justification for Evidence Category	<p>Sjögren’s syndrome associated neuropathy comprises a heterogeneous group of neuropathies. There is a very low level of evidence for IVIg use in this condition, with conflicting reports of efficacy. Improvement has been reported in small case series of patients with some forms of sensory neuropathy (Pereira 2016, Rist 2011, Yamashita 2013). Conflicting reports exist for patients with ataxic sensory neuronopathy, with one study reporting benefit (Takahashi 2003) but most studies (Rist 2011, Pereira 2016) and clinical experience suggesting it is ineffective for this patient group.</p> <p>As there may be a subset of patients who benefit from Ig therapy for this condition the Specialist Working Groups (Immunology and Neurology) of the Immunoglobulin Governance Program have recommended to retain this condition within the <i>Criteria</i>, but only where demonstrable benefit is shown. Where no demonstrable benefit can be identified, patients will not be eligible to remain on Ig therapy under these criteria.</p>
Diagnosis Requirements	A diagnosis must be made by a Neurologist.

Qualifying Criteria for Ig Therapy

Severe, primary Sjögren's syndrome associated neuropathy that is unresponsive to corticosteroid and immunosuppressant therapy

Use this indication for new patients and those that have never trialled off from Ig therapy. Please use the indication **Relapse of Sjögren's syndrome associated neuropathy within six months of trial off Ig therapy** for responding patients who have relapsed after weaning from Ig therapy.

- Primary Sjögren's syndrome associated neuropathy without necrotising vasculitis
- AND
- Significant disability due to neuropathy as measured by an [adapted Modified Rankin Scale \(MRS\) score](#) of at least three points

AND

- Unresponsive to an appropriate trial of corticosteroids
- OR
- Corticosteroids are contraindicated or have resulted in unacceptable side effects or significant toxicity

AND

- Unresponsive to at least one immunosuppressant agent
- OR
- Immunosuppressant medication is contraindicated

IVIg should be used for a maximum period of four months (Induction plus three maintenance cycles) before determining whether the patient has responded. If there is no benefit after this treatment, IVIg therapy should be abandoned.

Review by a neurologist is required within four months of treatment to determine whether the patient has responded. Continuing six monthly reviews may be undertaken by neurologists, rheumatologists or immunologists.

After a year of therapy a trial of Ig weaning should be attempted with a view to cessation within six months unless there is a contraindication to doing so.

Documentation of clinical effectiveness is necessary for continuation of IVIg therapy.

	<p>Relapse of Sjögren’s syndrome associated neuropathy within six months of trial off Ig therapy</p> <p>This indication should be used for responding Sjögren’s syndrome patients who have relapsed within six months of commencement of a trial off immunoglobulin therapy. For those who have not received Ig therapy previously please use the indication Severe, primary Sjögren’s syndrome associated neuropathy that is unresponsive to corticosteroid and immunosuppressant therapy</p> <ul style="list-style-type: none"> • Deterioration of proven neuropathy and in the adapted Modified Rankin Scale (MRS) score, of at least one point, in a previously stable patient as compared to the review score prior to the trial off Ig therapy <p>AND</p> <ul style="list-style-type: none"> • Relapse occurs within six months of the last immunoglobulin dose in a previously responding patient <p>IVIg should be used for a maximum period of four months (induction plus three maintenance cycles) before determining whether the patient has responded. If there is no benefit after this treatment, IVIg therapy should be abandoned.</p> <p>Review by a neurologist is required within four months of treatment to determine whether the patient has responded. Thereafter the continuing six monthly reviews may be undertaken by a neurologist, rheumatologist or immunologist.</p> <p>After a year of therapy a trial of Ig weaning should be attempted with a view to cessation within six months unless there is a contraindication to doing so.</p> <p>Documentation of clinical efficacy is necessary for continuation of IVIg therapy.</p> <p>Once a patient has relapsed when trialled off Ig treatment, a second line immunomodulatory agent should be strongly considered as additional therapy.</p>
Exclusion Criteria	Sjögren’s syndrome associated vasculitic neuropathy

Review Criteria for Assessing the Effectiveness of Ig Use

Severe, primary Sjögren’s syndrome associated neuropathy that is unresponsive to corticosteroid and immunosuppressant therapy

Review by a neurologist is required within four months (induction plus three maintenance cycles) of treatment to determine whether the patient has responded. If there is no benefit after this treatment, IVIg therapy should be abandoned. In responding patients, continuing six monthly reviews may be undertaken by neurologists, rheumatologists or immunologists.

After a year of therapy a trial of Ig weaning should be attempted with a view to cessation within six months unless there is a contraindication to doing so.

Documentation of clinical effectiveness is necessary for continuation of IVIg therapy.

Clinical effectiveness of Ig therapy can be assessed by:

On review of the initial authorisation period

- Improvement of neuropathy (formally assessed) compared to the qualifying assessment
AND
- Improvement (or no deterioration) in disability as measured by a decrease of at least one point (or no change) in the [adapted Modified Rankin Scale \(MRS\) score](#) compared to the qualifying assessment

On review of a continuing authorisation period

- Improvement in or stabilisation of neuropathy compared to the previous review assessment
AND
 - Improvement in or stabilisation of the degree of disability as measured by the [adapted Modified Rankin Scale \(MRS\) score](#) compared to the previous review assessment
- AND
- After 12 months on therapy a trial of Ig weaning towards cessation of Ig therapy is planned or a reason provided as to why a trial is not planned

Review by a neurologist is required within four months (induction plus three maintenance cycles) of treatment to determine whether the patient has responded. If there is no benefit after this treatment, IVIg therapy should be abandoned. In responding patients, continuing six monthly reviews may be undertaken by neurologists, rheumatologists or immunologists.

After a year of therapy a trial of Ig weaning should be attempted with a view to cessation within six months unless there is a contraindication to doing so.

Documentation of clinical effectiveness is necessary for continuation of IVIg therapy.

Once a patient has relapsed when trialled off Ig treatment, a second line immunomodulatory agent should be strongly considered as additional therapy.

Clinical effectiveness of Ig therapy may be assessed by:

On review of the initial authorisation period

- Improvement of neuropathy (formally assessed) compared to the qualifying assessment for this relapse indication
- AND
- Improvement in disability (or no further deterioration) as measured by a decrease of at least one point (or no change) in the [adapted Modified Rankin Scale \(MRS\) score](#) compared to the qualifying assessment

On review of a continuing authorisation period

- Improvement in or stabilisation of neuropathy compared to the previous review assessment
- AND
- Improvement in or stabilisation of the degree of disability as measured by the [adapted Modified Rankin Scale \(MRS\) score](#) compared to the previous review assessment
- AND
- After 12 months on therapy a trial of Ig weaning towards cessation of Ig therapy is planned or a reason provided as to why a trial is not planned

Once a patient has relapsed in the first six months of a trial off therapy, a further trial might be considered after at least one year.

Dose	<div data-bbox="518 33 1540 89">Severe, primary Sjögren's syndrome associated neuropathy that is unresponsive to corticosteroid and immunosuppressant therapy</div> <div data-bbox="582 145 1484 280"> <ul style="list-style-type: none"> • Induction Dose (IVIg) - 1- 2 g/kg in 2-5 divided doses • Maintenance Dose (IVIg) - 0.4-1g/kg, 4-6 weekly. The amount per dose should be titrated to the individual's response. A maximum dose of 1 g/kg may be given in any 4 week period. This might be small doses more frequently than fortnightly. </div> <div data-bbox="518 291 1540 353">The aim should be to use the lowest dose possible that achieves the appropriate clinical outcome for each patient.</div> <div data-bbox="518 380 1540 443">Refer to the current product information sheet for further information on dose, administration and contraindications.</div>
	<div data-bbox="518 510 1465 544">Relapse of Sjögren's syndrome associated neuropathy within six months of trial off Ig therapy</div> <div data-bbox="582 600 1500 734"> <ul style="list-style-type: none"> • Induction Dose (IVIg) - 1- 2 g/kg in 2-5 divided doses • Maintenance Dose (IVIg) - 0.4g/kg -1 g/kg, 4-6 weekly. The amount per dose should be titrated to the individual's response. A maximum dose of 1 g/kg may be given in any 4 week period. This might be small doses more frequently than fortnightly. </div> <div data-bbox="518 745 1540 808">The aim should be to use the lowest dose possible that achieves the appropriate clinical outcome for each patient.</div> <div data-bbox="518 835 1540 898">Refer to the current product information sheet for further information on dose, administration and contraindications.</div>

Bibliography

- Bonita, R & Beaglehole R 1988, 'Recovery of motor function after stroke' *Stroke*, vol. 19, no. 12, pp. 1497-500. <https://www.ncbi.nlm.nih.gov/pubmed/3201508>
- Brito-Zeron, P, Akasbi, M, Bosch, X, et al 2013, 'Classification and characterisation of peripheral neuropathies in 102 patients with primary Sjögren's syndrome', *Clinical and Experimental Rheumatology*, vol. 31, no. 1, pp. 103-10. <http://www.clinexprheumatol.org/abstract.asp?a=5938>
- Danieli, MG, Pettinari, L, Morariu, R, et al 2012, 'Intravenous immunoglobulin and mycophenolate mofetil for long-standing sensory neuropathy in Sjögren's syndrome', *Case reports in immunology*, Volume 2012, Article ID 186320, 3 pages. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4207592/>
- Molina, JA, Benito-Leon, J, Bermejo, F, et al 1996, 'Intravenous immunoglobulin therapy in sensory neuropathy associated with Sjögren's syndrome', *Journal of neurology, neurosurgery, and psychiatry*, vol. 60, no. 6, pp.699. <http://jnnp.bmj.com/content/60/6/699.1.short>
- Morozumi, S, Kawagahira Y, Iijima M, et al 2009, 'Intravenous immunoglobulin treatment for painful sensory neuropathy associated with Sjögren's syndrome', *Journal of the Neurological Sciences*, vol. 279, pp. 57-61. <https://www.ncbi.nlm.nih.gov/pubmed/19168191>
- Pereira, PR, Viala, K, Maisonnobe, T, et al 2016, 'Sjögren Sensory Neuropathy – Long-term outcome and treatment Response in a series of 13 cases', *Medicine (Baltimore)*, vol. 95, no. 19, pp. e3632. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4902517/>
- Rankin J 1957, 'Cerebral vascular accidents in patients over the age of 60', *Scottish medical journal*, vol. 2, no. 4, pp. 127-36.
- Rist, S, Sellam, J, Hachulla, E, et al 2011, 'Experience of intravenous immunoglobulin therapy in neuropathy associated with primary Sjögren's syndrome: A national, multicentric retrospective study', *Arthritis care & research*, vol. 63, no. 9, pp. 1339-44. <https://www.ncbi.nlm.nih.gov/pubmed/21584943>
- Stroke Engine Canada , 'The Modified Rankin Scale'. Available from [Modified Rankin Scale \(MRS\) – Strokengine](#)
- Takahashi, Y, Takata, T, Hoshino, M, et al 2003, 'Benefit of IVIg for long-standing ataxic sensory neuronopathy with Sjögren's syndrome', *Neurology*, vol. 60, no. 3, pp. 503-5. <https://www.ncbi.nlm.nih.gov/pubmed/12578938>
- Tzioufas, AG & Voulgarelis, M 2007, 'Update on Sjögren's syndrome autoimmune epithelitis: from classification to increased neoplasias', *Best Practice & Research Clinical Rheumatology*, vol. 21, issue. 6, pp. 989-1010. [http://www.bprclnrheum.com/article/S1521-6942\(07\)00105-2/fulltext](http://www.bprclnrheum.com/article/S1521-6942(07)00105-2/fulltext)
- Wakasugi, D, Kato, T, Gono, T, et al 2009, 'Extreme efficacy of intravenous immunoglobulin therapy for severe burning pain in a patient with small fiber neuropathy associated with primary Sjögren's syndrome', *Modern rheumatology*, vol. 19, no. 4, pp. 437-40. <https://www.ncbi.nlm.nih.gov/pubmed/19458906>
- Smith, AJ, Jackson, MW, Wang, F, et al 2005, 'Neutralisation of muscarinic receptor autoantibodies by intravenous immunoglobulin in Sjögren's syndrome', *Human Immunology*, vol. 66, no. 4, pp. 411–416. <https://www.sciencedirect.com/science/article/abs/pii/S0198885905000339>
- Van Swieten, JC, Koudstaal, PJ, Visser, MC, et al 1988, 'Interobserver agreement for the assessment of handicap in stroke patients', *Stroke*, vol. 19, no. 5, pp. 604-607. <https://www.ncbi.nlm.nih.gov/pubmed/3363593>

Bibliography

Yamashita, H, Eri, T, Ueda, Y, et al 2013, 'Diagnosis and treatment of primary Sjögren syndrome-associated peripheral neuropathy: a six-case series', *Modern Rheumatology*, vol. 23, no. 5, pp. 925-33. <http://link.springer.com/article/10.1007/s10165-012-0767-x>

Generated on: 30 November 2023